



## Steiner Tree:

- Input:
  - Graph  $G = (V, E)$
  - Costs  $c: E \rightarrow \mathbb{R}^+$
  - **Terminals**  $T \subseteq V$



- Feasible solutions:  $F \subseteq E$  s.t.  $F$  connected, spans all terminals
- Objective:  $\min_{F \subseteq E} \sum_{e \in F} c(e) = \min_F c(F)$

MST: Steiner tree with  $T = V$   
 shortest path: ST with  $T = \{s, t\}$



- terminals
- Steiner nodes (non-terminals)


Def:  $d: V \times V \rightarrow \mathbb{R}_{\geq 0}$  is a **metric space** on  $V$  if:

- $d(u, v) = 0$  iff  $u = v$
- $d(u, v) = d(v, u) \quad \forall u, v \in V$
- $d(u, v) \leq d(u, w) + d(w, v) \quad \forall u, v, w \in V$  (triangle inequality)



**Metric Steiner Tree:** (special case of ST on a metric space)

- Input:  $V$ , metric  $c: V \times V \rightarrow \mathbb{R}_{\geq 0}$  on  $V$ , terminals  $T \subseteq V$
- Feasible:  $F \subseteq V \times V$  s.t.  $F$  connected, spans all terminals
- Objective:  $\min \sum_{e \in F} c(e)$



Thm: If there is an  $\alpha$ -approx for Metric ST, then there is an  $\alpha$ -approx for Steiner Tree

Def: The **metric completion**  $c'$  of  $(G=(V,E),c)$  is the metric on  $V$  where  $c'(u,v)$  is the cost of the shortest path between  $u$  and  $v$  under edge lengths  $c$

Lemma: Let  $H$  be a solution (a Steiner Tree) for Steiner Tree problem on input  $(G,c,T)$ . Then  $H$  is a solution to Metric ST problem on input  $(V,c',T)$  with  $c'(H) \leq c(H)$


Pf:  $H$  feasible for metric: ✓

$c'(u,v) \leq c(u,v)$  by def of  $c'$   $\forall (u,v) \in E$

$$\Rightarrow c'(H) = \sum_{e \in H} c'(e) \leq \sum_{e \in H} c(e) = c(H) \quad \checkmark$$

Lemma: Let  $H'$  be a solution to Metric Steiner Tree on  $(V, c', T)$ . Then there is some solution  $H$  to Steiner Tree on  $(G, c, T)$  with  $c(H) \leq c'(H')$ , and given  $H'$  we can find  $H$  in polytime.

PF: Replace each  $\{u, v\} \in H'$  by shortest  $u-v$  path in  $G$   
 $\Rightarrow$  subgraph  $\hat{H}$  of  $G$ ,  $c(\hat{H}) \leq c'(H')$



Let  $H$  arbitrary spanning tree of  $\hat{H}$

$\Rightarrow c(H) \leq c(\hat{H}) \leq c'(H')$

PF of reduction thm:

Let  $A$   $\alpha$ -approx for Metric ST. Given input  $(G, c, T)$ , run  $A$  on  $(V, c', T)$  to get  $H'$ , use previous lemma to get  $H$ .

Let  $OPT_{\text{metric}}$  be opt solution for  $(V, c', T)$

$OPT$  be opt solution for  $(G, c, T)$

$$\begin{aligned}
c(H) &\leq c'(H') && \text{(lemma)} \\
&\leq \alpha \cdot c(\text{OPT}_{\text{metric}}) && \text{(def of } A) \\
&\leq \alpha \cdot c(\text{OPT}) && \text{(def of OPT}_{\text{metric}}) \\
&\leq \alpha \cdot c(\text{OPT}) && \text{(first lemma)}
\end{aligned}$$

So just need to design good alg for metric case

ALG:

- Return  $F = \text{MST on terminals!}$

Claim:  $F$  is valid solution

PF:  $F$  is: connected and spans  $T$



Def:  $G$  is **Eulerian** if there is a closed tour that uses every edge exactly once

Thm:  $G$  is Eulerian iff connected, all degrees even (even holds for multigraphs).

Thm: ALG is a  $2(1 - \frac{1}{|T|})$ -approximation

Pf:


Let  $F^*$  optimal solution.

WTS:  $c(F) \leq 2(1 - \frac{1}{|T|}) \cdot c(F^*)$


Plan: Find some spanning tree  $\hat{F}$  of  $T$  s.t.

$$c(\hat{F}) \leq 2(1 - \frac{1}{|T|}) \cdot c(F^*)$$

$\Rightarrow c(F) \leq c(\hat{F})$  since  $F$  MST of  $T$



start with  $F^*$



Double every edge:  $2F^*$

All degrees even: Eulerian!

Tour  $C$  which uses every edge:

$$c(C) = c(2F^*) = 2c(F^*)$$

"shortcut"  $C$  to only use terminals, see each terminal once: cycle  $H$

Triangle inequality:

$$c(H) \leq c(C)$$

Remove heaviest edge of  $H$ : path  $\hat{F}$

$$c(\hat{F}) \leq (1 - \frac{1}{|T|}) c(H) \leq 2(1 - \frac{1}{|T|}) c(F^*) \quad \checkmark$$

Metric TSP:

Input: Metric space  $(V, c)$

Feasible: Hamiltonian cycle  $H$

← cycle visiting all nodes once

Objective:  $\min c(H) = \sum_{e \in H} c(e)$

Alg 1:

- Compute MST  $T$
- Double  $T$  to get  $2T$
- $2T$  Eulerian, so Eulerian tour  $C$
- Shortest  $C$  to get  $H$

Thm:  $2(1 - \frac{1}{n})$  - approx

Prf: Just like Steiner Tree!

Let  $H^*$  optimal solution,

$F$  path from removing heaviest edge from  $H^*$

$$\Rightarrow c(H) \leq c(C) = c(2T) = 2c(T) \leq 2c(F)$$

shortest  $\nearrow$

$\nearrow$   
 $F$  a spanning tree

$$\leq 2(1 - \frac{1}{n})c(H^*)$$

Want to do better: Christofides' Algorithm

Why did we lose 2?

- Doubling MST

Why did we do that?

- Make it Eulerian

Cheaper way to make MST Eulerian?

Problem: odd degree nodes

Lemma: Let  $G=(V,E)$  be a graph. Then there are an even # nodes with odd degree.

Pf:

$$\sum_{v \in V} d(v) = 2|E| \quad (\text{even})$$



Def: A **perfect matching** of  $S \subseteq V$  is a matching on  $S$  of size  $\frac{|S|}{2}$  (every node in  $S$  matched to other node in  $S$ )



Fact: Can find min-cost perfect matchings in polytime

Christofides':

- Compute MST  $T$
- Let  $D$  be odd-degree nodes in  $T$
- Compute min-cost perfect matching  $M$  of  $D$
- Let  $C$  be Eulerian tour of  $T+M$
- Return  $H = \text{shortcutted } C$

Claim: Everything well-defined

Thm:  $\frac{3}{2}$ -approximation


PF: Let  $H^*$  optimal solution

$$c(T) \leq c(H^*)$$

$$c(H) \leq c(C) = c(T) + c(M) \leq c(H^*) + c(M)$$

$$\text{So wTS } c(M) \leq \frac{1}{2} c(H^*)$$

Start with  $H^*$  to  $D$ , get  $H_0$



$|D|$  even, so partition into "evens"  $M_1$  and "odds"  $M_2$   
- Each a perfect matching of  $D$

$$c(M_1) + c(M_2) = c(H_0)$$

$$c(M) \leq \min(c(M_1), c(M_2)) \leq \frac{1}{2} c(H_0) \leq \frac{1}{2} c(H^*)$$